

CONSTRUCTING AND DECONSTRUCTING GROUP ACTIONS

ALEJANDRO ADEM

ABSTRACT. Given a finite group G , it is not hard to show that it can act freely on a product of spheres. A more delicate issue is the following question: what is the minimum integer $k = k(G)$ such that G acts freely on a finite complex with the homotopy type of a product of k spheres? The study of this problem breaks up into two distinct aspects: (1) proving bounds on $k(G)$ in terms of subgroup data for G ; and (2) constructing explicit actions on a product of k spheres. In this note we will discuss aspects of both problems, including recent progress based on the periodicity methods developed in [5]. We also describe a potential counterexample to the prevalent expectations as well as recent work on constructing geometric actions on actual products of spheres.

1. INTRODUCTION

In 1944 P. A. Smith proved [24] that if a finite group G acts freely on a sphere, then all of its abelian subgroups must be cyclic. This condition is known to be equivalent to the *periodicity* of the cohomology of G with trivial integral coefficients—recall that the cohomology of G is said to be periodic if there exists a $d > 0$ such that $H^i(G, \mathbb{Z}) \cong H^{i+d}(G, \mathbb{Z})$ for all $i > 0$. These groups are characterized by having generalized quaternion 2-Sylow subgroups and cyclic p -Sylow subgroups for p odd.

In 1957, Milnor [20] showed that there are restrictions on G imposed by the geometry of the action: if G acts freely on a sphere $\mathbb{S}^n$, then every involution in G must be central. For example, the symmetric group on three letters Σ_3 cannot act freely on any sphere, even though it has periodic cohomology. In 1960 Swan [26] showed that a converse of Smith's result does hold for homotopy spheres; he proved that if G has periodic cohomology then it acts freely on a finite complex $X \simeq \mathbb{S}^n$ for some $n > 0$. After that there ensued substantial activity on the problem of spherical space forms, which culminated in 1976 with the solution due to Madsen, Thomas and Wall [19]; namely they proved that a finite group G acts freely on some sphere $\mathbb{S}^n$ if and only if all of its abelian subgroups are

Partially supported by the NSF. This paper is a modified version of the plenary lecture delivered by the author at the March 2002 Algebraic Topology Conference held at Northwestern University.

cyclic and every element of order 2 in G is central. In other words, the combination of Smith's homological condition and Milnor's geometric condition were enough to ensure the existence of a free action on a sphere.

Based on the above one can inquire about the analogous situation for groups which do not have periodic cohomology. In particular given such a group, can we construct a free action on a suitably *minimal* product of spheres? In this note we will survey aspects of this problem, and in particular describe recent substantial progress for groups which do not contain $\mathbb{Z}/p \times \mathbb{Z}/p \times \mathbb{Z}/p$ as a subgroup.

2. FREE ACTIONS ON A PRODUCT OF SPHERES

We can define the p -rank of a finite group G as $r_p(G) = \max \{r \mid (\mathbb{Z}/p)^r \subseteq G\}$ and its rank as $r(G) = \max \{r_p(G) \mid p \text{ divides } |G|\}$. Smith's pioneering work can be reformulated as saying that if a finite group G acts freely on a finite complex $X \simeq \mathbb{S}^n$, then $r(G) = 1$. Motivated by this and other evidence, the following rather difficult conjecture emerges:

Conjecture 2.1. If G acts freely on $X = \mathbb{S}^{n_1} \times \cdots \times \mathbb{S}^{n_m}$, then $r(G) \leq m$.

This question has been settled in the affirmative for $m = 2$ by Conner (in the equidimensional case [13]) and by Heller [18]. Better general results exist for an arbitrary product of equidimensional spheres; we recall a result due to Adem and Browder [2].

Theorem 2.2. *If G acts freely on a finite complex $X \simeq (\mathbb{S}^n)^k$, then $r_p(G) \leq k$ for p an odd prime. If $p = 2$, then this holds for $n \neq 1, 3, 7$.*

The homologically trivial case was completely settled by Carlsson [9], [10]. More recently the result above was extended to the case $p = 2$ and $n = 1$ by Yalcin [28]. The other cases remain intriguingly open. A geometrically interesting situation arises when the action of G permutes a basis for the homology of the product of spheres. Here we have a stronger bound, expressed in the following result, due to Adem and Benson [3].

Theorem 2.3. *Let G be an elementary abelian p -group of rank r acting freely on a finite dimensional CW complex $X \simeq (\mathbb{S}^n)^t$ in such a way that the basis $u_1, u_2, \dots, u_t$ of $H^n(X, \mathbb{F}_p)$ corresponding to the t spheres is permuted by G . Then the number of orbits of G on $\{u_1, \dots, u_t\}$ is at least r .*

From this we infer that if G is an elementary abelian p group acting freely on $X = (\mathbb{S}^n)^k$ and permuting the canonical basis, then $r(G) \leq \dim H_n(X, \mathbb{F}_p)^G$. Note that if $p = 2$ and $n \neq 1, 3, 7$, then every action satisfies this hypothesis (by a Hopf invariant

one argument, see [1]). We should also mention that Conjecture 2.1 follows from a more general conjecture (due to Carlsson, see [12]) arising from commutative algebra, namely:

Conjecture 2.4. If C_* is a free, finite and connected $\mathbb{F}_p G$ chain complex, where $G = (\mathbb{Z}/p)^r$, then $\sum_{i=0}^{\infty} \dim_{\mathbb{F}_p} H_i(C) \geq 2^r$.

This has been verified in some instances; when $p = 2$ it is known to hold for $r_2(G) \leq 4$ (see [11] and [23]); this was also settled in the general case of 1-dimensional chain complexes in [6].

Methods from the cohomology of groups are used to study these problems. To illustrate some of the techniques we will prove a simple yet useful result which can help us approach Conjecture 2.1.¹ Recall that BG denotes the classifying space of a group G ; given any free G -action X , there is a classifying map $X/G \rightarrow BG$ associated to the action.

Proposition 2.5. *Let $G = (\mathbb{Z}/2)^r$ act freely on a closed n -manifold M such that the classifying map $\pi_G : M/G \rightarrow BG$ induces a surjection onto the top non-zero mod 2 cohomology group $H^n(M/G, \mathbb{F}_2)$. Then the action does not extend to a free action of a larger elementary abelian 2-group.*

Proof. Assume then that G acts as stated in the theorem, but that the action extends to a free action by an elementary abelian 2-group K , with $[K : G] = 2$. Recall that given an index two subgroup there is a Gysin sequence associated to it in equivariant cohomology, which is natural with respect to equivariant maps. Hence we obtain a diagram of Gysin sequences, where we assume coefficients are in $\mathbb{F}_2$:

$$\begin{array}{ccccc} H^n(BG) & \xrightarrow{\text{tr}} & H^n(BK) & \xrightarrow{\cup\chi} & H^{n+1}(BK) \\ \downarrow \pi_G^* & & \downarrow \pi_K^* & & \downarrow \pi_K^* \\ H^n(M/G) & \xrightarrow{\text{tr}} & H^n(M/K) & \xrightarrow{\cup\chi} & H^{n+1}(M/K) \end{array}$$

Here $\chi \in H^1(BK)$ is the cohomology class associated to the index two subgroup $G \subset K$. Choose $0 \neq x \in H^n(M/K)$, then as $H^{n+1}(M/K) = 0$, there exists a class $0 \neq x' \in H^n(M/G)$ such that $x = \text{tr}(x')$. Using our hypotheses, we can find an $\alpha \in H^n(BG)$ with $\text{tr}(\pi_G^*(\alpha)) = x$. However by the commutativity of the diagram, we have that this expression is the same as $\pi_K^* \text{tr}(\alpha) = 0$, as the transfer map is zero for elementary abelian 2-groups. The result follows. $\square$

¹Although this result was known to the author for some time, it should be attributed to [16], where this method has been explained in full detail.

For example, one can show that the hypotheses of this theorem hold for actions of $\mathbb{Z}/2 \times \mathbb{Z}/2$ on a product of two spheres (see [16] for details) and more generally for actions of $(\mathbb{Z}/2)^r$ on $X = (\mathbb{S}^n)^r$ (see [9]).

From this we derive a conjecture which would imply Conjecture 2.1, namely:

Conjecture 2.6. If $G = (\mathbb{Z}/2)^r$ acts freely on $X = \mathbb{S}^{n_1} \times \cdots \times \mathbb{S}^{n_m}$, then $\pi_G^* : H^N(G, \mathbb{F}_2) \rightarrow H^N(X/G, \mathbb{F}_2)$ is surjective, where $N = n_1 + \cdots + n_m$.

Remark 2.7. The case of $(\mathbb{Z}/2)^r$ actions on a product of real projective spaces is also of some interest. For free actions, the main result there (see [7]) is: if $(\mathbb{Z}/2)^r$ acts freely on a finite complex $X \simeq (\mathbb{R}P^n)^m$, then

$$r \leq \begin{cases} 0 & n \equiv 0, 2 \pmod{4} \\ m & n \equiv 1 \pmod{4} \\ 2m & n \equiv 3 \pmod{4} \end{cases}$$

The case of non-free actions is worthy of some attention. For example, we have that if $(\mathbb{Z}/2)^k$ acts on $X = (\mathbb{S}^m)^t$, then there exists an $x \in X$ with $\dim G_x \geq k - t$ (this follows from the methods used in [9]). In contrast, actions on real projective spaces behave very differently; for example we have (see [7]):

Proposition 2.8. *There exists an action of $G = (\mathbb{Z}/2)^{1249}$ on $X = (\mathbb{R}P^{2^{1298}-1})^{50}$, with isotropy subgroups of rank less than or equal to 50.*

This example arises from representations for certain class 2 nilpotent groups, introduced by Ol'shanskii [21].

3. CONSTRUCTING FREE ACTIONS

Given the results in the previous section, a natural problem is that of *constructing* free actions on a product of $r(G)$ spheres. In his landmark paper [26], Swan showed that if $r(G) = 1$, then G in fact acts freely on a finite complex $X \simeq \mathbb{S}^n$. The natural extension of this result leads to a difficult open problem:

Conjecture 3.1. A finite group G acts freely on a finite complex X which has the homotopy type of a product of $r(G)$ spheres.

This conjecture is due to Benson and Carlson. In a purely algebraic context, this was completely settled in [8]. The geometric side of the problem is very difficult, as it amounts to explicitly constructing free group actions on products of spheres.

The most basic examples of spheres with actions of finite groups arise from representation theory. Given a unitary or orthogonal G -representation V , the unit vectors $S(V)$ are homeomorphic to a sphere. In many cases one can choose representations $V_1, V_2, \dots, V_r$ such that G acts freely on $S(V_1) \times \dots \times S(V_r)$. However, there are some restrictions, as the following result due to U. Ray ([22]) shows.

Proposition 3.2. *If G is a finite group acting freely on a product of spheres arising from representations, then the only possible non-abelian composition factors of G are the alternating groups A_5 and A_6 .*

Example 3.3. Let P denote the p -Sylow subgroup of $\mathrm{GL}_3(\mathbb{F}_p)$; it is an extra-special group of order p^3 . It is not hard to show that P acts freely on a product $S(V_1) \times S(V_2) \times S(V_3)$; however more is true (see [22]): in fact the *minimal* number of representations required for a free action on a product is always three.

Example 3.4. In [21], it was shown that the alternating group A_4 cannot act freely on any finite complex $X \simeq \mathbb{S}^n \times \mathbb{S}^n$, for any n . Hence in particular A_4 cannot act freely on any $\mathbb{S}^n \times \mathbb{S}^m$ via a product action, as we could produce a free action on a product of two equidimensional spheres by taking the appropriate joins.

To construct new examples of free group actions we will build on what we know about groups with periodic cohomology. In fact we will consider G -CW complexes such that the isotropy subgroups all satisfy this condition. The following result appears in [5].

Theorem 3.5. *Let X denote a simply-connected finite G -CW complex such that all of its isotropy subgroups have periodic cohomology. Then there exists a finite complex $Y \simeq X \times \mathbb{S}^n$ with a free G -action.*

This result can be proved directly (see [27]) by constructing a spherical fibration over successive skeleta in X . This uses coherence provided by a “universal” periodicity class $\alpha \in H^*(G, \mathbb{Z})$ for actions with periodic isotropy; classical spherical space forms and the fact that all obstructions can be killed using fiber joins. This is based on the methods first developed in [14]. The result is also a consequence of a more general theorem in [6] about spaces with periodic cohomology. This notion (requiring twisted coefficients) is induced by cup product with a given cohomology class.

From the previous result we see that a clear new strategy emerges for constructing free group actions: to get a free G -action on $X \times \mathbb{S}^n$ for some $n > 0$, it suffices to build an action

of G on X with periodic isotropy.² We now look more carefully at finite group actions on spheres; given $X \simeq \mathbb{S}^N$ with a G -action, we have an Euler class $\alpha \in H^{N+1}(G, \mathbb{Z})$ associated to it.

Definition 3.6. An Euler class $\alpha \in H^{N+1}(G, \mathbb{Z})$ is said to be effective if $H^*(X \times_G EG, \mathbb{Z})$ has Krull dimension less than the rank $r(G)$.

Question 3.7. Given a finite group G , does there exist an effective Euler class in $H^*(G, \mathbb{Z})$?

If we assume X to be finite dimensional, then we see that the Euler class α is effective if and only if $X^E = \emptyset$ for all $E \subset G$ elementary abelian subgroups of maximal rank.

We now focus on the case of a finite p -group P . It follows from known results (see [17]) that if an effective Euler class exists, then an effective Euler class must arise from a representation sphere $S(V)$. Let $C \cong \mathbb{Z}/p$ be a central subgroup in P , and let $V = \text{Ind}_C^P(\chi)$ where χ is a non-trivial character for C . Then $X = S(V)$ has an action of P such that C acts freely. Hence $X^E = \emptyset$ for all E elementary abelian subgroups of maximal rank and so we obtain an effective Euler class $\alpha \in H^{2[P:C]}(P, \mathbb{Z})$. More generally, if the center $Z \subset P$ has rank equal to z , then we can find representations $V_1, V_2, \dots, V_z$ of P such that Z acts freely on the product $S(V_1) \times \dots \times S(V_z)$. Note that the isotropy for the action of P will have rank equal to $r(P) - r(Z)$. This indicates that for any p -group we can construct at least part of the desired free action on a product of spheres “for free”. This is analogous to the fact that we always have a regular sequence of length $r(Z)$ in $H^*(P, \mathbb{F}_p)$.

4. RANK TWO GROUPS

We now consider groups G with $r(G) = 2$. If P is a p -group of rank equal to two, then we know that it will act on a sphere $X = S(V)$ with periodic isotropy. Hence we obtain (see [5]):

Theorem 4.1. *A finite p -group P acts freely on a finite complex $X \simeq \mathbb{S}^n \times \mathbb{S}^m$ if and only if it does not contain a subgroup isomorphic to $(\mathbb{Z}/p)^3$.*

Of course in particular this establishes Conjecture 3.1 for $r(P) = 2$. In addition the conjecture has been settled affirmatively for almost all rank 2 simple groups. Indeed, we have

²These techniques also apply to infinite groups, and they can be used to show (see [5]) that a discrete group Γ acts freely and properly discontinuously on $\mathbb{S}^n \times \mathbb{R}^m$ for some $m, n > 0$ if and only if Γ is countable and has periodic cohomology.

Theorem 4.2. *If G is a finite simple group with $r(G) = 2$ other than $PSL_3(\mathbb{F}_p)$, then it acts freely on a finite complex $X \simeq \mathbb{S}^n \times \mathbb{S}^m$; moreover any such action must be exotic, i.e. it cannot be a product action.*

The proof that such an action cannot be a product action is a consequence of the classification of finite simple groups! Indeed, one checks every non-abelian simple group contains a copy of A_4 , and we know that this group cannot act freely via a product action.

Example 4.3. Let $G = SL_3(\mathbb{F}_2)$, $|G| = 2^3 \cdot 3 \cdot 7$. From the character tables in the *Atlas of Finite Groups* [15], we note that there exists a 3-dimensional complex representation V of G such that $V^E = \{0\}$ for every $E = \mathbb{Z}/2 \times \mathbb{Z}/2$ in G . Hence $X = S(V)$ is a G -space with periodic isotropy, and so, applying our previous results, G acts freely on a finite complex $Y \simeq \mathbb{S}^5 \times \mathbb{S}^m$ for some m . However in this case we can proceed more directly as follows. The subgroup $G \subset U(3)$ must actually lie in $SU(3)$, as it is a simple group. Then if we consider the $SU(2)$ -fibration $SU(3) \rightarrow SU(3)/SU(2) = \mathbb{S}^5$, we see that it is G -equivariant and that the G -action on the total space is free. Hence our group acts freely on the total space E of an $SU(2)$ -bundle over $\mathbb{S}^5$, which we can regard as the sphere bundle of a 2-dimensional complex G -vector bundle τ over $\mathbb{S}^5$. This corresponds (non-equivariantly) to the non-zero element in $\pi_4(SU(2)) \cong \mathbb{Z}/2$. Now the sphere bundle $S(\tau \oplus \tau)$ is a trivial bundle, as any $\mathbb{S}^7$ -bundle over $\mathbb{S}^5$ splits as a product (see [25], page 139). Hence $G = SL_3(\mathbb{F}_2)$ acts freely and smoothly on the manifold³ $M = \mathbb{S}^5 \times \mathbb{S}^7$.

Our techniques reduce the problem of constructing a free action on a product of two spheres to constructing an action on a single sphere with periodic isotropy. Equivalently we can construct an action on a sphere such that the associated Euler class is effective. Although this is a considerable reduction, there are still groups which cannot be handled with this approach. In particular we have a result due to Unlu [27]:

Proposition 4.4. *If $G = PSL_3(\mathbb{F}_p)$ acts on a finite complex $X \simeq \mathbb{S}^n$, then there exists a subgroup $H \cong \mathbb{Z}/p \times \mathbb{Z}/p$ in G such that $X^H \neq \emptyset$.*

In fact this result has been recently generalized by Grodal and Smith to show that if G acts on *any* homotopy sphere X , then the equivariant cohomology $H^*(EG \times_G X, \mathbb{F}_p)$ has Krull Dimension equal to two, hence is not periodic in the sense of [5] and so there is no hope of constructing a free G action on a product of two spheres using the methods

³More generally we have shown that if G is any finite subgroup of $SU(3)$, then it acts freely and smoothly on $\mathbb{S}^5 \times \mathbb{S}^7$; this is a special case of a result in [4].

above. In particular this gives an example of a group for which Question 3.7 has a negative answer.

Motivated by these considerations and in view of some very special properties of the group cohomology $H^*(PSL_3(\mathbb{F}_3), \mathbb{F}_3)$, it seems valid to raise the following

Question 4.5. Does $G = PSL_3(\mathbb{F}_3)$ in fact act freely on a finite complex $X \simeq \mathbb{S}^n \times \mathbb{S}^m$?

Needless to say it would be rather significant if this question had a negative answer.

On the other hand this seems to be a rather special situation, which does not apply for example to odd order groups. Work in progress (joint with Grodal) indicates that every odd order group of rank equal to two will indeed act freely on a finite $X \simeq \mathbb{S}^n \times \mathbb{S}^m$. This requires a detailed analysis of their p -Sylow subgroups and fusion properties.

5. GEOMETRIC ACTIONS

So far we have not discussed the more geometric side of constructing group actions. Ideally we would like a group of rank equal to r to act freely on an actual product of r spheres—by this we mean a manifold homeomorphic to it, or even better, diffeomorphic. Although this has been settled in the rank one case, very little is currently known for higher rank groups. Given our previous results in the homotopy category, a good question to raise would be

Question 5.1. If P is a p -group with $r(P) = 2$, does it act freely on the manifold $M = \mathbb{S}^n \times \mathbb{S}^m$ for some $m, n > 0$?

This problem is rather difficult as it naturally leads into both homotopy-theoretic and surgery related issues. We can however offer a very recent result in this direction (joint work with Davis and Unlu [4]):

Theorem 5.2. *Let $G \subset U(n)$ be a faithful unitary representation of G , such that the induced action of G on $U(n)/U(k)$ is free. Then, if $|G|$ is relatively prime to $(n-1)!$, the group G will act freely on $M = \mathbb{S}^{2n-1} \times \dots \times \mathbb{S}^{2k+1}$.*

Corollary 5.3. *Let P denote a p -group with cyclic center and an abelian maximal subgroup. Then P acts freely on $M = \mathbb{S}^{2p-1} \times \dots \times \mathbb{S}^{2p-r-1}$ where $r < p$ is the rank of P .*

Example 5.4. We consider the extra-special p -group P of order p^3 and exponent p . Applying the above, it will act freely on $M = \mathbb{S}^{2p-1} \times \mathbb{S}^{2p-3}$.

The proof of the theorem above requires an interesting combination of group theory, homotopy theory and surgery theory. The basic strategy is to propagate the free action on the Stiefel manifold to a product of spheres. The corollary follows from the classification of those finite p -groups which admit a faithful unitary representation in $U(p)$ such that the induced action on the complex Stiefel manifold $U(p)/U(p-r)$ is free.

REFERENCES

- [1] Adem, A., $\mathbb{Z}/p$ Actions on $(\mathbb{S}^n)^k$, Transactions of the AMS **300** (1987), pp. 791–809.
- [2] Adem, A. and Browder, W., *The Free Rank of Symmetry of $(\mathbb{S}^n)^k$* , Inventiones Mathematicae **92** (1988), pp. 431–440.
- [3] Adem, A. and Benson, D.J., *Elementary Abelian Groups Acting on Products of Spheres*, Math. Zeitschrift **228** (1998), pp. 705–712.
- [4] Adem, A., Davis, J. and Unlu, O., *Fixity and Group Actions*, in preparation.
- [5] Adem, A. and Smith, J., *Periodic Complexes and Group Actions*, Annals of Mathematics **154** (2001), pp. 407–435.
- [6] Adem, A. and Swan, R.G., *Linear Maps Over Abelian Group Algebras*, J.Pure and Applied Algebra **104** (1995), pp. 1–7.
- [7] Adem, A. and Yalcin, E., *On Some Examples of Group Actions and Group Extensions*, J.Group Theory **2** (1999), pp. 69–79.
- [8] Benson, D. and Carlson, J., *Complexity and Multiple Complexes*, Math. Zeitschrift **195** (1987), pp. 221–238.
- [9] Carlsson, G., *On the Nonexistence of Free Actions of Elementary Abelian Groups on Products of Spheres*, Amer. Journal Math. **102** (1980), pp. 1147–1157.
- [10] Carlsson, G., *On the Rank of Abelian Groups Acting Freely on $(\mathbb{S}^n)^k$* , Inventiones Mathematicae **69** (1982), pp. 393–400.
- [11] Carlsson, G., *Free $(\mathbb{Z}/2)^3$ Actions on Finite Complexes*, Annals of Math. Studies **113** (1987), pp. 332–344.
- [12] Carlsson, G., *Free $(\mathbb{Z}/2)^k$ Actions and a Problem in Commutative Algebra*, Lecture Notes in Math **1217** (1985), pp. 79–83.
- [13] Conner, P., *On the Action of a Finite Group on $\mathbb{S}^n \times \mathbb{S}^n$* , Annals of Math. **66** (1957), pp. 586–588.
- [14] Connolly, F. and Prassidis, S., *Groups Which Act Freely on $\mathbb{R}^m \times \mathbb{S}^{n-1}$* , Topology **28** (1989), pp. 133–148.
- [15] Conway, J. et al., **Atlas of Finite Groups**, Oxford University Press, Eynsham (1985).
- [16] Dammak, M., Grazzini, F. and Zarati, S., *Actions Libres d’un 2-Groupe Abélien Elementaire Sur un Produit Fini de Spheres*, preprint 2001.
- [17] Dotzel, R. and Hamrick, G., *p -group Actions on Homology Spheres*, Inv.Math. **62** (1981), pp. 437–442.
- [18] Heller, A., *A Note on Spaces with Operators*, Ill. J. Math. **3** (1959), pp. 98–100.

- [19] Madsen, I., Thomas, C.B. and Wall, C.T.C., *The Topological Spherical Space Form Problem II*, Topology **15** (1978), pp. 375–382.
- [20] Milnor, J., *Groups Which Act on $\mathbb{S}^n$ without Fixed Points*, Amer.J.Math. **79** (1957), pp. 623–630.
- [21] O’lshanskii, A.Yu., *The Number of Generators and Orders of Abelian Subgroups of Finite p -groups*, Math.Notes **23** (1978), pp. 183–185.
- [22] Ray, U., *Free Linear Actions of Finite Groups on Products of Spheres*, J.Algebra **147**, pp. 456–490.
- [23] Refai, M., *Group Actions on Finite CW Complexes*, Indian J.Pure and Applied Math. **24** (1993), pp. 245–255.
- [24] Smith, P.A., *Permutable Periodic Transformations*, Proc.Nat.Acad.Sci. **30** (1944), pp. 105–108.
- [25] Steenrod, N., **The Topology of Fibre Bundles**, Princeton University Press 1951.
- [26] Swan, R.G., *Periodic Resolutions for Finite Groups*, Annals of Mathematics **72** (1960), pp. 267–291.
- [27] Unlu, O., *Doctoral Dissertation, UW-Madison*, in preparation.
- [28] Yalcin, E., *Group Actions and Group Extensions*, Transactions AMS **352** (2000), pp. 2689–2700.

MATHEMATICS DEPARTMENT, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN, 53706

E-mail address: adem@math.wisc.edu